

Case 2804

***Drosophila putrida* Sturtevant, 1916 (Insecta, Diptera): proposed replacement of the holotype by a neotype**

David Grimaldi

*Department of Entomology, American Museum of Natural History,
Central Park West at 79th Street, New York, N.Y. 10024-5192, U.S.A.*

Abstract. The purpose of this application is to designate a neotype in accordance with current usage for the nominal species *Drosophila putrida* Sturtevant, 1916. Examination of the holotype shows that it belongs to an un-named species which has been consistently misidentified as *Drosophila testacea* von Roser, 1840. *D. putrida* is widely used in ecological, genetic and evolutionary studies and is restricted to the eastern U.S.A.

1. North America has two species belonging to the small, Holarctic *Drosophila testacea* species group. Their species status and nomenclature have never been critically examined and some confusion exists, partly as a result of the long and consistent misidentification of *Drosophila putrida* Sturtevant, 1916. The holotype of *putrida* is a male in perfect condition in the American Museum of Natural History (type locality: Woods Hole, Massachusetts, U.S.A.). I recently examined the holotype and its paratypes. The specimens actually belong to a species which since about 1940 has been misidentified as *Drosophila testacea* von Roser, 1840.

2. Confusion began when two externally distinct North American species in the group were fully recognized, and the name *testacea* was applied to the species most similar to the true (European) *testacea*, although current work has shown that the North American '*testacea*' is a different, morphocryptic species. Few voucher specimens exist in collections from all the biological work done on the two North American species, so it is impossible to confirm the identity of the *putrida*/*'testacea'* referred to in older papers. However, there are specimens collected in Austin, Texas in the 1940's in the University of Texas collection at the American Museum of Natural History which have labels identifying *putrida* in the sense recognized today. Patterson & Stone (1952) distinguished the two species on the basis still adhered to, as does Strickberger's (1962) key which is in wide use today. Apparently, no one had ever checked Sturtevant's type specimen of *putrida*.

3. The three species in the *testacea*-group are abundant inhabitants of forests, and have been favored subjects for studies in ecology, genetics and evolution. An extensive literature exists; major papers that treat either one or both of the Nearctic species are the following: Carson & Stalker, 1951 (breeding sites); Dorsey & Carson, 1956 (host finding behavior); Grimaldi, 1985 (niche biology); Grimaldi & Jaenike, 1983 (*putrida* hosts), 1984 (larval competition); Jaenike, 1978, 1986 (host selection), 1988 (parasitism of '*testacea*'); Jaenike & Grimaldi, 1983 (oviposition population genetics); Jaenike et al., 1983 (toxin resistance); James & Jaenike, 1990 ('sex ratio' meiotic drive); Lacy, 1982, 1983, 1984 (host use and population genetics); Levitan, 1954 (distributional

records); Miller & Weeks, 1964 (distributional records); Montague & Jaenike, 1985 (parasitism); Patterson & Stone, 1952 (distributions, internal reproductive organs, distinguishing characters, chromosomes); Patterson & Wagner, 1943 (distributions); Patterson & Wheeler, 1949 (North American *Drosophila* catalogue); Sabath, Richmond & Torella, 1973 (temperature controlled color polymorphism); Strickberger, 1962 (key to North American *Drosophila*); Throckmorton, 1962a, 1962b, 1975 (*Drosophila* phylogeny); Ward, 1949 (metaphase chromosomes); Wharton, 1943 (metaphase chromosomes); Wheeler, 1981a (world catalogue); Wheeler, 1981b (Nearctic fauna). Adoption of *putrida* in the sense of the holotype would cause serious confusion because the name, as used in the above literature, would be transferred to the other species. The references listed in this paragraph all agree upon a diagnosis of *putrida* as having a pair of presutural acrostichal setulae that are stouter, decumbent and only about twice the length of other, standard acrostichal setulae.

4. A revision of the *testacea*-group is completed, utilizing adult specimens from all known localities of the range, as well as electrophoresis studies, mating tests and ecological characteristics. There is no doubt that the species represented by the *D. putrida* neotype proposed below, from New Jersey, also occurs in the locality (Massachusetts) of the holotype and that no other species share the diagnostic traits of the proposed neotype.

5. In accordance with Recommendation 75E of the Code, I refer the case to the Commission to set aside the existing type material of *D. putrida* and to confirm the designation of a neotype belonging to the taxonomic species that North American *Drosophila* workers have been consistently referring to as *putrida* for the last 50 years. What has been called '*testacea*' in North America needs a new name, diagnosis and designated type. The *putrida* neotype I propose is an adult male specimen labelled as '*Drosophila* (*D.*) *putrida* Sturtevant, 1916, NEOTYPE, Det. D.A. Grimaldi' from 'U.S.A.: New Jersey: Morris County, Pompton Plains, June, 1986, D.A. Grimaldi, coll.', and deposited in the American Museum of Natural History. No problem would exist in reconciling Sturtevant's original (1916) and subsequent (1921) descriptions of *putrida* with the neotype, since he omitted crucial diagnostic details of the presutural setae which externally distinguish the species. His description could apply to any of the *testacea*-group species; indeed, it was this insufficiently detailed description that contributed to the continued misidentification.

6. The International Commission on Zoological Nomenclature is accordingly asked:

- (1) to use its plenary powers to set aside all previous fixations of type specimens for the nominal species *Drosophila putrida* Sturtevant, 1916 and to confirm the neotype designation proposed in para. 5 above;
- (2) to place on the Official List of Specific Names in Zoology the name *putrida* Sturtevant, 1916, as published in the binomen *Drosophila putrida* and as defined by the neotype designated in (1) above.

Acknowledgements

I am grateful to Curtis W. Sabrosky for his suggestions on an early draft of this proposal, and to John Jaenike and Avis James (University of Rochester) for additional references and collaborative work on the species status of American and European '*testacea*'.

References

- Carson, H.L. & Stalker, H.D. 1951. Natural breeding sites for some wild species of *Drosophila* in the Eastern United States. *Ecology*, **32**: 317–330.
- Dorsey, C.K. & Carson, H.L. 1956. Selective responses of wild Drosophilidae to natural and artificial attractants. *Annals of the Entomological Society of America*, **49**: 177–181.
- Grimaldi, D. 1985. Niche separation and competitive coexistence in mycophagous *Drosophila* (Diptera: Drosophilidae). *Proceedings of the Entomological Society of Washington*, **87**: 498–511.
- Grimaldi, D. & Jaenike, J. 1983. The Diptera breeding on skunk cabbage, *Symplocarpus foetidus* (Araceae). *Journal of the New York Entomological Society*, **91**: 83–89.
- Grimaldi, D. & Jaenike, J. 1984. Competition in natural populations of mycophagous *Drosophila*. *Ecology*, **65**: 1113–1120.
- Jaenike, J. 1978. Host selection by mycophagous *Drosophila*. *Ecology*, **59**: 1286–1288.
- Jaenike, J. 1986. Intraspecific variation for resource use in *Drosophila*. *Biological Journal of the Linnean Society*, **27**: 47–56.
- Jaenike, J. 1988. Parasitism and male mating success in *Drosophila testacea*. *American Naturalist*, **131**: 774–780.
- Jaenike, J. & Grimaldi, D. 1983. Genetic variation for host preference within and among populations of *Drosophila tripunctata*. *Evolution*, **37**: 1023–1033.
- Jaenike, J., Grimaldi, D.A., Sluder, A.E. & Greenleaf, A.L. 1983. α -Amanitin tolerance in mycophagous *Drosophila*. *Science*, **221**: 165–167.
- James, A.C. & Jaenike, J. 1990. 'Sex ratio' meiotic drive in *Drosophila testacea*. *Genetics*, **126**: 651–656.
- Lacy, R.C. 1982. Niche breadth and abundance as determinants of genetic variation in populations of mycophagous drosophilid flies (Diptera: Drosophilidae). *Evolution*, **36**: 1265–1275.
- Lacy, R.C. 1983. Structure of genetic variation within and between populations of mycophagous *Drosophila*. *Genetics*, **104**: 81–94.
- Lacy, R.C. 1984. Predictability, toxicity, and trophic niche breadth in fungus-feeding Drosophilidae (Diptera). *Ecological Entomology*, **9**: 43–54.
- Leviton, M. 1954. Drosophilidae in New York and New Jersey. *The American Midland Naturalist*, **52**: 453–459.
- Miller, D.D. & Weeks, L. 1964. *Drosophila* collections near the Blue Ridge of southwestern North Carolina. *The American Midland Naturalist*, **72**: 93–114.
- Montague, J.R. & Jaenike, J. 1985. Nematode parasitism in natural populations of mycophagous drosophilids. *Ecology*, **66**: 624–626.
- Patterson, J.T. & Stone, W.S. 1952. *Evolution in the genus Drosophila*. 610 pp. MacMillan, New York.
- Patterson, J.T. & Wagner, R.P. 1943. Geographical distribution of species of the genus *Drosophila* in the United States and Mexico. *University of Texas Publications*, **4313**: 217–281.
- Patterson, J.T. & Wheeler, M.R. 1949. Catalogue of described species belonging to the genus *Drosophila*, with observations on their geographical distribution. *University of Texas Publications*, **4920**: 207–233.
- Sabath, M.D., Richmond, R.C. & Torella, R.M. 1973. Temperature-mediated seasonal color changes in *Drosophila putrida*. *The American Midland Naturalist*, **90**: 509–512.
- Strickberger, M.W. 1962. Key to United States species of the genus *Drosophila*. Pp. 111–122 in Strickberger, M.W. (Ed.), *Experiments in Genetics with Drosophila*. Wiley, New York.
- Sturtevant, A.H. 1916. Notes on North American Drosophilidae with descriptions of twenty-three new species. *Annals of the Entomological Society of America*, **9**: 323–343.
- Sturtevant, A.H. 1921. The North American species of *Drosophila*. *Carnegie Institute of Washington Publications*, **301**: 1–150.
- Throckmorton, L.H. 1962. The problem of phylogeny in the genus *Drosophila*. *University of Texas Publications*, **6205**: 207–343.

- Throckmorton, L.H.** 1962. The use of biochemical characteristics for the study of problems of taxonomy and evolution in the genus *Drosophila*. *University of Texas Publications*, **6205**: 415–487.
- Throckmorton, L.H.** 1975. The phylogeny, ecology, and geography of *Drosophila*. Pp. 421–469 in King, R.C. (Ed.), *Handbook of genetics*, vol. 3. Plenum, New York.
- Ward, C.L.** 1949. Karyotype variation in *Drosophila*. *University of Texas Publications*, **4920**: 70–79.
- Wharton, L.T.** 1943. Analysis of the metaphase and salivary chromosome morphology within the genus *Drosophila*. *University of Texas Publications*, **4313**: 282–319.
- Wheeler, M.R.** 1981a. The Drosophilidae: a taxonomic overview. Pp. 1–97 in Ashburner, M., Carson, H.L. & Thompson, J.N. (Eds.), *The genetics and biology of Drosophila*, vol. 3a. 429 pp. Academic Press, New York.
- Wheeler, M.R.** 1981b. Geographical survey of Drosophilidae: Nearctic species. Pp. 99–121 in Ashburner, M., Carson, H.L. & Thompson, J.N. (Eds.), *The genetics and biology of Drosophila*, vol. 3a. 429 pp. Academic Press, New York.